# FOR MORE INFORMATION

www.mylymphedema.com

**National Lymphedema Network** 

www.lymphnet.org 1-800-541-3259

**Lymphedema Awareness Foundation** 

www.eLymphNodes.org 407-324-3255

**Lymphovenous Society of Canada** 

www.lymphovenous-canada.ca

Y-ME National Breast Cancer Organization

www.y-me.org 1-800-221-2141 (24-Hour patient hotline)

**Healthy Steps** 

www.gohealthysteps.com 877-365-6014

1-888-255-1300 www.juzousa.com











Since 1912

Visit www.juzousa.com for more information.



What You Should Know About


LYMPHEDEMA



**Juzo**<sup>®</sup>

Since 1912

Copyright © 2013 by Julius Zorn, Inc. All rights reserved, especially the right to reproduce and translate into foreign languages. No part of this brochure may be reproduced in any way without prior written consent of Julius Zorn Inc. DBA Juzo. ® Juzo is a registered trademark of Julius Zorn, Inc. The Lebed Method is a registered trademark of Focus on Healing Inc.




The following information is for people who have had mastectomies, lumpectomies, radiation treatment, lymph node removal, surgeries, infections, or trauma to the limbs. You may be at risk for developing lymphedema. Or perhaps you or a family member were diagnosed at birth, during childhood or early adulthood with lymphedema. This information will help you understand what lymphedema is and how it can be managed.

"One day I awoke and my arm and hand were swollen.


I thought I would just elevate it for awhile and the swelling would subside. Days later and much to my frustration, nothing happened. Little did I know, this was not going to be a temporary thing. What could it be? What caused this? What was I going to do? I want you to read on, knowing this information really saved my life."

—E.D.



# IF YOU KNOW SOMEONE WHO MAY BE AT RISK FOR DEVELOPING LYMPHEDEMA:

Please be sure he or she reads this brochure and gives a copy to his or her doctor. Creating awareness and understanding about lymphedema is our goal.



# How Does LYMPHEDEMA DEVELOP?

Lymphedema most often develops in one arm or leg, but may be present in both arms and both legs. It may also occur in the hands or feet, and even in the chest, back, neck, face, abdomen, and genitals.



## PRIMARY LYMPHEDEMA

Primary Lymphedema is caused by malformations of the lymphatic system. These malformations are most common in women. Symptoms may be present at birth or may develop later, often during puberty or pregnancy. Primary lymphedema is most common in the legs, but may also occur in the arms or torso.

## SECONDARY LYMPHEDEMA

Secondary Lymphedema is a result of damage to the lymphatic system. Surgical procedures such as mastectomies, lumpectomies with radiation and/or removal of lymph nodes are the most common causes. Secondary lymphedema occurs most commonly in the arms, but may also develop in the legs. Other causes may include a traumatic injury, infection, or severe chronic venous insufficiency.



## WHAT ARE THE SYMPTOMS?

Lymphedema may develop within a few months after a procedure, years later, or not at all. The first obvious sign of lymphedema is swelling characterized by "pitting." This is recognizable when the skin is depressed for a few seconds and the indentation does not immediately disappear. Other symptoms may include a tightness or heaviness in the affected area or changes in the texture of the skin. You may even notice that jewelry and clothing feel tighter..



## WHAT TO DO IF SYMPTOMS OCCUR?

If your lymphatic system is compromised, this area of the body will have to work much harder to circulate fluid. When it is unable to keep up, swelling and fluid build-up are likely to occur. That is why the lymphatic vessels in the compromised area need external support.

## WEARING A COMPRESSION GARMENT

Wearing a compression garment provides external pressure to assist lymph drainage and hopefully prevent additional swelling. Although swelling is sometimes reversible, in most cases it is not and could lead to chronic lymphedema.

When you are about to engage in repetitive motions, such as exercise, cleaning, yard work, golfing or tennis it is a good idea to wear a compression garment. It is also important to wear a compression garment when you are flying. The decrease in cabin pressure in the airplane puts stress on the lymphatic and circulatory systems and this can increase swelling.

WHAT IS THE TREATMENT FOR LYMPHEDEMA?

Once the diagnosis of lymphedema is confirmed, certain treatment procedures are indicated. Since there is no cure for lymphedema, the goal of

are indicated. Since there is no cure for lymphedema, the goal of treatment is to reduce the swelling and to maintain the reduction. For a majority of patients, this can be achieved by the skillful application of Complete Decongestive Therapy (CDT) which is safe, reliable, and non-invasive. It is proven to provide positive long-term results for both primary and secondary lymphedema. CDT is performed by specially trained therapists. The treatment consists of four components:

- Manual Lymph Drainage (MLD)
- Graduated Compression Garments or bandages
- Meticulous Skin Care
- Therapeutic Exercises



lymph fluid out of the affected region and reduce the swelling using MLD techniques and bandaging. Ideally these treatments are performed daily, five days a week for 2-4 weeks, depending on the severity. Once the swelling is reduced, the patient is fitted with a graduated compression garment. This marks the second phase, in which the patient plays a large role in his or her ongoing self-care. Wearing graduated compression garments, being meticulous about skin care, and performing self-manual lymph drainage and therapeutic exercises will assure continued success.

Treatment procedures such as diuretics or surgery are not appropriate for lymphedema. Initially, a diuretic will decrease the water content and reduce swelling, but the protein molecules remain in the tissues and swelling will reoccur as soon as the drug loses its effectiveness. Surgical methods for lymphedema have not proven successful.





# A CLOSER LOOK AT THE TREATMENT

### 1. Manual Lymph Drainage (MLD):

Manual Lymph Drainage is necessary to manage fluid volume of the affected extremity to a normal or near normal size. Bandages are used in the phase 1 and compression garments in phase 2.

### **2.** Graduated Compression Garments:

Graduated compression garments are necessary to maintain the reduced limb and are designed to replace the bandages that were used earlier in treatment. They are worn during the day while you are active. At night when you sleep, bandages or night garments are usually worn. The compression garments help to keep the swelling down, improve circulation, and prevent the reaccumulation of lymph fluid in the area. The garments must be worn every day, for life.

### **3.** Pneumatic Compression Pumps:

Some treatment may include use of a pneumatic compression pump. This mechanical device works as a sleeve with chambers that are intermittently inflated with air. This device is sometimes used in combination with CDT.

### 4. Skin Care:

With lymphedema, the skin is usually dry and may crack easily, making it very susceptible to infections. A low-pH lotion, free of alcohol and fragrances should be used to maintain the moisture of the skin and to protect it. If an infection develops, consult your physician immediately!

### 5. Exercise:

A skilled therapist may recommend an exercise program that is suited for your particular need and ability. An effective decongestive exercise program aids the muscular system to move lymph fluid out of the affected area. Any exercise that causes pain or muscle soreness should be discussed with a therapist. Moderate exercise such as swimming, walking, biking, light weight training and yoga are recommended. Overly aggressive sports that may cause injury should be avoided. Consult your physician before beginning any program.



## LIFESTYLE

# TIPS FOR MANAGING AND LIVING WITH LYMPHEDEMA


If you are at risk for developing lymphedema or already have it, these guidelines will help you prevent and manage the condition. Although you may not need to follow each suggestion, it is good to be aware of them.

If you have lymphedema, it is important that you avoid injury and overexertion in the affected area. You should wear the prescribed compression garments and/or bandages as instructed by your physician or therapist. Be sure all of your medical charts are updated and consider wearing a lymphedema alert bracelet available through the National Lymphedema Network.



# 1. CLOTHING & JEWELRY

Avoid tight or restrictive clothing that may impair circulation or cause irritation and swelling. Shirt sleeves, bra straps, belts, socks, and shoelaces should fit comfortably. Shoes with low heels are recommended. Jewelry, including watches, bracelets, and rings should not feel tight. Avoid carrying heavy purses, luggage and shoulder bags on the side that is affected.



### 2. SKIN CARE

Keep skin protected using a moisturizing lotion and wash with a mild pH balanced soap. When bathing, water should be warm, not hot. Avoid irritating or allergy-causing cosmetics, detergents, deodorants, and perfumes. Inspect your feet, hands, and limbs regularly for skin changes since open skin invites bacteria and infection. Avoid cuts, pricks, scratches, burns, and insect bites that could increase swelling or promote infection. Use sunscreen and insect repellent regularly. Gloves are also a good protective measure for the hands when washing dishes, cooking, cleaning or working outdoors.



# TRAUMA CAN ALSO INCREASE SWELLING & FURTHER DAMAGE THE LYMPHATIC SYSTEM

Avoid blood pressure checks, IV's, blood draws, and shots in the affected limb. Do not cut into the cuticles when trimming fingernails or toenails. Use an electric razor when shaving.

## 3. PROPER NUTRITION

Increased weight complicates lymphedema. Maintain a balanced diet of fruit, vegatables, whole grains and protien. Avoid excessive consumption of fatty foods, sweets, salt and alcohol. Drink plenty of water and unsweetened liquids.

## 4. EXERCISE

Moderate exercise such as swimming, walking, biking, light strength training and yoga are recommended. Overly aggressive sports that may cause injury should be avoided. Consult your physician before beginning any program.

# 5. Weather & Traveling

Always wear compression garments when flying or driving, as recommended by your physician or therapist. When traveling by car, be sure the seatbelt is comfortable and make frequent stops since prolonged sitting may increase swelling. Avoid extreme changes in temperature (over 90 degrees or below zero), sun bathing, tanning beds, saunas, steam rooms, and hot tubs.

# **FAQS**

### WILL LYMPHEDEMA EVER GO AWAY?

If it is diagnosed early, lymphedema can be managed through proper treatment, quality compression garments, and good skin care. That is why management is key.

77777777777777777777777777777777

### WHERE DO I FIND AN MLD THERAPIST?

Many hospitals have trained MLD therapists in their physical and occupational therapy department or outpatient rehabilitation centers. Lymphedema organizations list MLD therapists on their websites. NALEA (North American Education Services) www.lymphedemaeducationassociation.org

#### WHERE CAN I BUY COMPRESSION GARMENTS?

Compression garments are sold at medical supply stores and some pharmacies. Measurements are taken by a certified fitter to ensure garments fit properly. Go to juzousa.com to find a local dealer.

# WILL MY INSURANCE PAY FOR COMPRESSION GARMENTS?

That depends on what type of insurance you have and what state you live in. Call your insurance company to verify coverage.

# HOW DO I CARE FOR MY COMPRESSION GARMENTS?

Daily machine wash and dry with a mild detergent is recommended to maximize therapeutic effectiveness. Do not use fabric softeners, chlorine or bleach which may damage your garment. Refer to the manufacturer's care instructions.

#### HOW DO I PUT COMPRESSION GARMENTS ON?

Wearing donning gloves is recommended to protect your garments from fingernail and jewelry snags. There are also numerous assistive devices that make putting on and taking off garments easy and comfortable. Ask your certified fitter for a demonstration of these devices.

### **GARMENTS**

Juzo offers a complete line of compression products for Lymphedema management. From compression stockings to arm sleeves, and hand gauntlets, Juzo garments are comfortable, durable and fashionable. They provide theraputic compression and have many added value features and benefits. Garments will maintain their compression for six months.







"I bought one pair to try and fell in love...and can't stop bragging about them and showing them off. Within one week of wearing them my legs felt and looked better! It is a wonderful feeling to be 'normal' again!"

- HRBS

"After breast cancer, I was told that I had lymphedema and the best garments to wear are Juzo, that I would enjoy the comfort, fit, and flexibility. And after wearing ten sleeves and gloves in the last three years, I am truly happy. Juzo has made lymphedema much more manageable. Thank-you"

- Brenda D.

"I am using your arm sleeve for two years with excellent results. I think your product is very comfortable and makes my life so easy! Thanks"

- Maria J

"I have found with lymphedema, that my patients have had more success and comfort with Juzo as compared to (competitors). I, as a physical therapist, have been very satisfied with the product and product line."

-Jamie D., Physical Therapist

For an easy approach to putting on and taking off compression garments, Juzo also offers the innovative Slippies System.

For a supplier of Juzo lymphedema garments near you, please call the Juzo Patient Care Line at **1-888-255-1300** or **juzousa.com**.