

NLN

NATIONAL LYMPHEDEMA NETWORK

LYMPH Link

Vol.19, No.1 January-March 2007

ISSN 1065-9951

Editor: EMMA VICTORIA G. BLANCO

Managing Editor: SASKIA R.J. THIADENS, RN

Graphics: MARCIA KRUSE
Cathedral City, CA

Editorial Committee: JANE ARMER, RN, PHD
MARCIA BECK, RN, CS, CMLDT
KATHLEEN FRANCIS, MD
BONNIE B. LASINSKI, MA, PT
CATHERINE M. TUPPO, PT

NATIONAL LYMPHEDEMA NETWORK
Latham Square
1611 Telegraph Avenue, Suite 1111
Oakland, CA 94612-2138
Telephone 510-208-3200
Website www.lymphnet.org
E-mail nin@lymphnet.org

The NLN Mission Statement

The mission of the NLN is to create awareness of lymphedema through education and to promote and support the availability of quality medical treatment for all individuals at risk for or affected by lymphedema.

The NLN is dedicated to:

- promoting research into the causes, prevention and treatment of lymphedema;
- securing adequate insurance coverage for medically necessary, safe and effective treatment;
- expanding the number and geographical distribution of lymphedema treatment facilities and certified therapists.

To achieve these goals, the NLN disseminates information about lymphedema to health care professionals so they can appropriately counsel their patients on its avoidance, and prescribe safe, effective treatment for those affected by this condition. The NLN also provides this information to the general public.

Understanding Lipedema

Guenter Klose, CI, CLT-LANA, and Roman H.K. Strößenreuther, MD

The term "lipedema" was first used in 1974 by ALLEN and HINES. Their publication is regarded as "the classical description" of the syndrome^{5, p. 184}:

"We wish to describe a clinical syndrome, lipedema of the legs, which is frequently very distressing. In our experience it affects solely women. The chief complaint is of swelling of the legs and feet...On questioning, the physician may elicit that enlargement of the limbs has always been generalized and symmetrical. The swelling below the knees is accentuated when patients are on their feet much and in warm weather. Aching distress in the legs is common...Occasionally, a patient feels, that her large legs have 'ruined her life.' Many are 'ashamed' of their legs."

This describes the patient's problems very well, but it is difficult to define lipedema precisely because the definition depends mainly on subjective findings. There are no medical or laboratory tests to distinguish local lipohypertrophy (local fat tissue increase) of the legs or hips from lipedema or general obesity. Consequently, lipedema is not generally accepted as a real disease. We will attempt to describe the current knowledge on the pathophysiology and treatment of lipedema.

Lipedema is a metabolic disorder of the adipose (fat) tissue with unknown etiology, affecting almost exclusively females. The following clinical and pathophysiological findings are of importance for differential diagnosis.

The term "lipedema" was chosen by ALLEN and HINES^{5,128} to describe a symmetrical swelling of both legs, extending from the hips to the ankles and excluding the feet, caused by an abnormal amount of subcutaneous adipose tissue. Typically, bulging masses can be found in the proximal thigh region and at the medial aspect of the knees. Occasionally, large, overhanging and hypersensitive fat lobes develop in these areas^{5,22,32,33,36,38}. In others, contours of the legs are more funnel-shaped, with a decrease of the adipose tissue noted below the knees. Some individuals present with similar changes in the arms especially when patient is also obese^{33,38,95}.

Most patients report slow onset of symptoms, often in connection with beginning of puberty^{33,35,36,49,86}; in other cases, there is no specific time of onset^{5,36,42,46,128}. Some patients report that other females in the family also suffer from lipedema^{42,59,132}. ALLEN and HINES¹²⁸ found a positive family history in 20% (n=119), but epidemiologic studies concerning

lipedema do not yet exist. There is no evidence of a specific genetic disorder or incidence related to race¹²⁸.




FIGURE 1:
Typical distribution pattern of the increased fat masses: 1=hip cushion; 2=proximal thigh bulge; 3=medial knee bulge; 4=fatty cuff at the malleoli; 5=local pre- or retromalleolar lipoma²².

Numerous impairments result from the abnormal pathophysiology of the tissues. The epidermis and subcutaneous tissues of the lipedematous

legs show a decreased elasticity^{5,42,66,67,128}. Aging causes further loss of skin elasticity^{18,80} and progression of the condition^{5,38,41,42,66,67,128}. Problems with normal ambulation (walking) lead to secondary orthopaedic deformities of the knees and feet^{16,17,22,33,36,38,42}. The tissue resistance to the contracting calf muscles (calf muscle pump) is too low and results in passive hyperaemia and an increase infiltration of water through blood capillaries, resulting in increased lymphatic water load. Swelling occurs when this fluid load exceeds the transport capacity of the lymphatic system^{42,66,67,80}. Blood capillaries are fragile and even insignificant trauma (e.g., hitting the leg on a table corner) can cause the development of small haematomas and a further increase in lymphatic load^{5,33,42,59,128}. Blood

Continued on page 2

In This Issue

PRESIDENT'S MESSAGE	4
CASE STUDY: FUNCTIONAL GAINS	5
7th NLN CONFERENCE REVIEW	7
CASE STUDY: LIPEDEMA	9
LEGISLATION UPDATE	11
QUESTION CORNER	12
RESOURCE GUIDE - 8-page pull-out	13
SUPPORT GROUPS	20
NEWS & NOTES	21
SCHOLARSHIP RECIPIENTS	26
EDUCATION CORNER	28
BECOME AN NLN SUPPORTER	31

Understanding...

Continued from cover page

coagulation tests for this condition are typically normal. It is not surprising that many patients develop an emotional disorder considering the physical appearance of the legs ALLEN and HINES pointed out:

"Patients with lipedema ordinarily are very sensitive about the appearance of their legs; they wear long skirts, avoid appearance in swimming suits and stand behind chairs at parties ... They are likely to be mirror peepers, searching repeatedly in mirrors for evidence that the appearance of their legs is not actually as bad as it seems to be"¹²⁸; p.124⁵. Evidence of neurosis is likely to be found. Occasionally, a patient feels that her large legs have 'ruined her life.' Many are 'ashamed' of their legs"¹⁰; p.55."

One patient described her situation:

"When you become older, you also become a little more overweight and your legs 'go through the roof.' You develop constant rubbing between your thighs; in summer it is itching and becomes sore. With age you also become more lazy—and, because of this constant itching and pain, you don't see any possibility for doing exercises. As the obesity worsens, lower back pain develops, the joints begin to hurt and your friends make silly jokes about your shape—and finally you rest at home, your only consolation is the chocolate in your nightstand and you will be unable to get out of this vicious circle!"

After some years, in cases of coexisting general obesity, lipedema can transform into lipo-lymphedema (STEMMER'S sign is now positive) in which fatty tissues begin to hold fluid¹⁴³. Furthermore, there is a correlation between obesity and disturbances of vascular edema protective reflexes^{9,24,25,51,52,53,54,60,65,79,85,100,101}, which leads to additional lymphatic water load. Edema is always the result of lymphatic insufficiency and, indeed, the lymphatic system shows typical pathological changes. The pre-lymphatic channels are widened; lymphatic capillaries have aneurysm-like changes⁶² and we find an irregular corkscrew-like course of the lymphatic vessels^{42,107,117}. The increased adipose tissue and the enlarged adipocytes most likely deform these thin lymphatic structures^{42,123}.

When lipo-lymphedema is present, pitting (indentable) edema in the lower legs can be noted when the person is upright or walking. This edema causes an uncomfortable feeling of tension. A reversal of the edema is possible after elevation

of the legs for a longer period or overnight^{5,41,42,49,127}.

Scintigraphy of the lymphatic system^{93,116,117,118,124} shows contradictory results. Sometimes, a reduced outflow in comparison to normal persons is described^{17,117,123} and other times, normal results were found^{21,124}. The decreased transport capacity of the lymphatics due to aging develops faster than in healthy legs⁴².

Unfortunately, there are no universally accepted guidelines for diagnosis of lipedema and, in some cases, the differentiation between lipedema and lipo-lymphedema is difficult^{47,119}. The typical progression of lipedema is as follows:

- In the **first stage** of the disease, the skin of the legs appears soft, smooth and regular. But palpation reveals changed structures of the adipose tissue that have a resemblance to "small Styrofoam balls"¹¹².
- In the **second and third stage**, the tissue structure becomes more and more nodular and tough, developing large deforming fatty lobes^{35,123}, especially at the inner side of the thighs, the knees and above the ankles^{22,42}.


DIFFERENTIAL DIAGNOSIS

The diagnosis of lipedema can be difficult in the early stage or if a combination form exists. Differential diagnosis of LE and lipedema is, in most cases, possible by taking the medical history and evaluation of the STEMMER'S sign. Distinction between primary bilateral lower extremity LE and lipedema can be difficult, although bilateral LE usually presents asymmetrically in contrast to symmetrical presentation of lipedema^{4,22,42,89,94,113}.

TREATMENT OF LIPEDEMA

While some of the pathophysiological tissue changes contributing to lipedema

FIGURE 2:



Typical shape of uni- and bilateral lymphedema and lipedema (l to r).


are identified, the real cause of the disease remains unknown. Therefore, therapy is predominantly symptom oriented. The goal must be to improve the disturbed lymph transport, the pathophysiological changes in microcirculation in the enlarged adipose tissue and a reduction of pain and the fat masses.

The most common therapeutic intervention for lipedema is Complete Decongestive Therapy (CDT)^{34,37,42,126,127}. The main constituents of this therapy concept are Manual Lymph Drainage (MLD)^{144,433} and compression therapy^{5,15,42,99,108,128}. Diet, skin care and remedial exercises are also very important.

At the *Lymphologica 1999* in Marburg/Germany, BRENKE reported, a volume reduction of 3.2 liters after a 3-week intensive therapy (mean initial volume of 23.3 liters; the average volume reduction at the thighs was 2.85 liters). In our Department of Lymphology (Freiburg/Germany), the mean volume reduction after 2 weeks of CDT is 14 % on average.

After initial decongestion through MLD and compression bandages, the patients are fitted with custom-made compression stockings. Permanent compression therapy causes significant reduction of adipose tissue^{15,35}, has a positive influence on the disturbed veno-arteriolar response and improves relative insufficiency of the

DIFFERENTIATION	LIPEDEMA	BILATERAL PRIMARY LYMPHEDEMA
Sex	almost exclusively women	more women than men
Beginning	often with the menarch	often with the menarche
Development	simultaneous begins at the whole legs	usually distal beginning
Extent	from the iliac crest to the ankle; no involvement of the dorsum of the feet	whole leg, involvement of the dorsum the feet
Stemmer's sign	negative	positive
Distribution	symmetric	asymmetric
Pain/hypersensitivity	yes	no
Skin temperature	decreased	normal
Skin color	normal, sometimes pale or cyanotic	normal
Haematoma	yes, even after minor injury	no
Tissue consistence	for a long time very soft tissue; later development of fibrosclerosis	progressive lymphostatic fibrosclerosis
Edema	pitting edema of the lower legs only after prolonged orthostasis	pitting in stage I, later fibrosclerosis
Dorsum of the feet	no edema	edema in most cases
Hyperkeratosis	no	yes
Cellulitis	no	often
Influence of positioning on edema	only decreases the orthostatic edema	significant decrease only in state I



Left 2 photos: Patient with lipo-lymphedema; before treatment and after 4 months of in-patient Phase-1 CDT given within one year. Weight reduction was 60 kg and leg volume reduction of 21 liters. Right 2 photos: Same patient, posterior view.

and improves relative insufficiency of the venous pump of the lower legs.

Exercise and sport activities are recommended, but the effect on adipolysis or the loss of leg and hip fat is often disappointing. Nevertheless, some younger patients report an improvement when working out 4-5 times a week, a minimum of 45 minutes and with an exertion rate of 75% (related to the maximum heart frequency). This recommendation may be unrealistic for some women. Some patients feel a positive effect after KNEIPP (hydrotherapy) applications; currently there is no research available concerning this therapy method.

Weight reduction is absolutely essential if patients are overweight because lipo-

lymphedema may develop particularly in patients who are also obese³⁶. There are, however, no specific dietary recommendations for lipedema.

Additional therapy with external pneumatic compression is sometimes recommended¹²⁶. Some patients report positive results with pneumatic compression, but controlled studies do not exist. ALLEN und HINES⁵ and others consider the use of diuretics a mistake.

In some select cases, liposuction has been recommended for the treatment of lipedema. Surgeons have shown good results, but problems also exist. After liposuction, some patients develop chronic lymphedema^{40,123,127}, lymphatic cysts or large haematoma⁶³ and have

problems with wound healing. Moreover, the cosmetic results are not always satisfactory⁶³. In recent years, the technique of liposuction has improved, and complications are fewer. Experience with a small number of patients indicates that the combination of surgical techniques with CDT may help to improve the results of liposuction. (Schmeller et al: Dtsch Arztebl 2005; 102:A 1061-1067 [Heft 15]). No definitive studies are available to confirm this finding.

CONCLUSION

Lipedema must be differentiated from local lipohypertrophy, primary LE of both legs and general obesity. Until the real cause of lipedema is known, treatment is symptomatic. With adequate treatment and optimal patient adherence, good results can be achieved, progression of lipedema can be halted and additional health problems prevented.

Guenter Klose, CI, CLT-LANA
Klose Training & Consulting, LLC,
Boulder, CO—guenter@klosetraining.com

Roman Strößenreuther, MD
Bavaria, Germany
stroessenreuther@t-online.de

References available through the NLN office or at www.lymphnet.org.

SECOND BIENNIAL NATIONAL LYMPHEDEMA NETWORK

International Patient Summit

OCTOBER 5-7, 2007

Lymphedema: Sharing Our World Of Knowledge

FOR PATIENTS, THOSE AT RISK, FAMILY MEMBERS, CAREGIVERS, ADVOCATES & THE GENERAL PUBLIC

Registration Available in Mid-January 2007 at www.lymphnet.org

Premier Sponsorship Opportunities Available ❖ Exhibitors Welcome ❖ Call 510-208-3200

Email: 2007patientsummit@lymphnet.org

RENAISSANCE WAVERLY HOTEL, ATLANTA, GEORGIA