

References

1. Stanton AW, Mellor RH, Cook GJ, et al. Impairment of lymph drainage in subfascial compartment of forearm in breast cancer related lymphedema. *Lymphat Res Biol.* 2003;1(2):121-132.
2. Finegold DN, Schacht V, Kimak MA, et al. HGF and MET mutations in primary and secondary lymphedema. *Lymphat Res Biol.* 2008;6(2):65-68.
3. Newman B, Lose F, Kedda MA, et al. Possible genetic predisposition to lymphedema after breast cancer. *Lymphat Res Bio.* 2012;10(1):2-13.
4. Kilbreath SL, Breast cancer. In Lee B-B, Bergan J, Rockson, SG, eds. *Lymphedema: A precise compendium of theory and practice.* London, UK: Springer. 2011:501-505.
5. Cemal Y, Pusic A, Mehrara BJ. Preventative Measures for Lymphedema: Separating Fact From Fiction. *J Am Coll Surg.* 2011;213(4):543-551.
6. National Lymphedema Network. Position Paper on Diagnosis and Treatment of Lymphedema. <http://www.lymphnet.org/pdfDocs/nlntreatment.pdf>. Updated Feb 2011.
7. Werner RS, McCormick B, Petrek J, et al. Arm edema in conservatively managed breast cancer: obesity is a major predictive factor. *Radiology.* 1991;180(1):177-184.
8. Helyer L, Varnic M, Le LW, Leonq W, McCreedy D. Obesity is a risk factor for developing post operative lymphedema in breast cancer patients. *Breast J.* 2010;16(1):48-54.
9. Ridner SH, Dietrich MS, Stewart BR, Armer JM. Body mass index and breast cancer-related lymphedema. *Support Care Cancer.* 2011;19(6):853-857.
10. Shaw C, Mortimer P, Judd PA. A randomized controlled trial of weight reduction as a treatment for breast cancer related lymphedema. *Cancer.* 2007;110(8):1868-1874.
11. National Lymphedema Network. Position Paper on Exercise. <http://www.lymphnet.org/pdfDocs/nlnexercise.pdf>. Updated Apr 2011.
12. Schmitz KH, Ahmed RL, Troxel A, et al. Weight lifting in women with breast-cancer-related lymphedema. *N Engl JM.* 2009;361(7):664-673.
13. Gavish I, Brenner B. Air travel and the risk of thromboembolism. *Intern Emerg Med.* 2011;6(2):113-116.
14. Feldman J. The Challenge of Infection in Lymphedema. *Lymphlink.* National Lymphedema Network. 2011. 23(3).
15. Ryan M, Stainton MC, Jaconelli C, Watts S, MacKenzie P, Mansberg T. The experience of lower limb lymphedema for women after treatment for gynecologic cancer. *Onc Nurs Forum.* 2003;30(3):417-423.
16. Vignes S, Dupuy A. Recurrence of lymphoedema-associated cellulitis (erysipelas) under prophylactic antibiotherapy: a retrospective cohort study. *J Eur Acad Dermatol Venereol.* 2006;20(7):818-822.
17. Mortimer PS. Swollen lower limb-2: lymphoedema. *BMJ.* 2000;320(7248):1527-1529.
18. Hettrick H, Nof L, Ward S, Ecthernach J. Incidence and prevalence of lymphedema in patients following burn injury: a five-year retrospective and three-month prospective study. *Lymphat Res Biol.* 2004;2(1):11-24.
19. Maheshwari M, Khandelwal S, Maheshwari S. Elephantiasis nostras: complications of third degree acid burn. *J Assoc of Physicians of India.* 2006;54:713.
20. Cole T. Risks and benefits of needle use in patients after axillary surgery. *Br JNurs.* 2006;15(18):969-979.
21. National Lymphedema Network. Position Paper on Screening and Measurement for Breast- Cancer-Related Lymphedema (BCRL). <http://www.lymphnet.org/pdfDocs/nlnBCRL.pdf>. Updated Feb 2011.
22. Schmitz KH, Ahmed RL, Troxel AB, et al. Weight lifting for women at risk for breast cancer-related lymphedema: a randomized trial. *JAMA.* 2010;304(24):2699-2705.
23. Ward LC, Battersby KJ, Kilbreath SL. Airplane travel and lymphedema: a case study. *Lymphology.* 2009;42(3):139-145.
24. Kilbreath SL, Ward LC, Lane K, et al. Effect of air travel on lymphedema risk in women with history of breast cancer. *Breast Cancer Res Treat.* 2010;120(3):649-654.
25. Graham PH. Compression prophylaxis may increase the potential for flight-associated lymphoedema after breast cancer treatment. *Breast.* 2002;11(1):66-71.
26. Greene AK, Borud L, Slavin SA. Blood pressure monitoring and venipuncture in the lymphedematous extremity. (letter to the editor) *Plast Reconstruc Surg.* 2005;116(7):2058-2059.
27. Zhang TS, Huang WY, Han LY, Liu WY. Heat and bandage treatment for chronic lymphedema of extremities. Report of 1045 patients. *Chin Med J (Engl).* 1984;97(8):567-577.
28. Chang TS, Han LY, Gan JL, Huang WY. Microwave: an alternative to electric heating in the treatment of peripheral edema. *Lymphology.* 1989;22(1):20-24.
29. Liu NF, Olszewski. The influence of local hyperthermia on lymphedema and lymphedematous skin of the human leg. *Lymphology.* 1993;26(1):28-37.

©2011 National Lymphedema Network (NLN). Permission to duplicate this handout as-is, in its entirety, for the educational purposes only, not for sale. All other rights reserved. For reprint permission, please contact the NLN at nln@lymphnet.org.